## Quarterly Report of Analytical Results for the CEMP Air Sampling Network

The CEMP air-sampling network is designed to monitor and collect radioactive airborne particles from NTS and non-NTS related activities, as well as background environmental sources. This report is provided to the station managers as a summary of the results from the analysis of the air samples they have collected as part of the community environmental monitoring program.

In general, the CEMP air-sampling network is comprised of 29 continuously operating environmental sampling stations. A total of 27 stations are equipped with a low volume air sampler/totalizer configuration to collect particulate radionuclides on glass fiber filter paper. Ideally, the samples are collected on a weekly basis with a target collection time of 168 hours (one week). The samplers are calibrated on a monthly basis by DRI to maintain a collection rate of 2.0 cfm (@ STP). All relevant information such as collection times, variations in flow rate, actual flow volumes, power outages, and other information documenting the integrity of the sample are recorded by the station managers. This allows for the proper interpretation of the analytical results. The air filters are analyzed by a commercial laboratory for gross alpha/beta activity as well as by high-resolution gamma spectrometry. The filters are composited on a quarterly basis (13 weeks) for gamma spectroscopy analysis only after the gross alpha/beta analyses have been completed.

The principle reporting units used in the U.S. for the measurement of radioactivity in the atmospheric environment is pCi/m<sup>3</sup> (picocuries per cubic meter). DRI receives its data from the lab as microcuries per filter, which is then recalculated to microcuries per milliliter based on the information provided by the station managers as well as monthly calibration results. This is the notation used for DRI internal databases and annual reports to DOE/NNSA. For the ease in constructing the tables contained in this report, as well as to hopefully the ease of comparison among stations and previous results, the units of pCi/m<sup>3</sup> are used. The data for the environmental TLD is reported in milliroentgens (mR).

A summary of the first quarter CY2007 analytical results for gross alpha and beta analyses are found in Tables 1 and 2. These tables show the minimum, maximum, and average values for each of the stations of the air-sampling network. The last column shows the average annual value from the previous year (CY2006) for comparison purposes. Overall the gross alpha results for the first quarter of CY2007 reflect similar values to the previous quarters. These data remain consistent with the average CY2006 analyses used for comparison, especially when analytical error is considered. The first quarter CY2007 beta results are also consistent with previous results.

The first quarter gamma results for CY2007 are shown in Tables 3. All of the samples are again gamma spectrum negligible (i.e. gamma emitting radionuclides were not detected) with the exception of Beryllium (Be)-7 and occasionally Lead (Pb)-210, both naturally occurring elements of the atmospheric and geological environment. Overall, these data are consistent with previous analytical results.

The TLD results for the first quarter of CY2007 are shown in Table 4. Overall, the results display similar values to the previous quarters of this calendar year. The 2006 PIC

exposure rate and TLD data are also shown for comparison. As with historical data, TLD values are commonly lower than the PIC results. The overall estimated annual exposure based on the first quarter shows consistent agreement with CY2006.

Finally, as station managers, your input concerning the contents of these reports are welcome and encouraged. We are interested in anything you feel would be helpful for you to interpret the data or to enable you to explain the information to someone in your community not familiar with the program.

(Average analytical error, +/- 0.0007)					
Station	Minimum (pCi/m <sup>3</sup> )	Maximum (pCi/m <sup>3</sup> )	Average (pCi/m <sup>3</sup> )	2006 Average (pCi/m <sup>3</sup> )	
Alamo	0.0009	0.0037	0.0021	0.0016	
Amargosa	0.0006	0.0018	0.0012	0.0012	
Beatty	0.0005	0.0045	0.0018	0.0013	
Boulder City	0.0011	0.0051	0.0029	0.0023	
Caliente	0.0005	0.0030	0.0021	0.0017	
Cedar City	0.0006	0.0019	0.0011	0.0011	
Delta	0.0005	0.0029	0.0015	0.0011	
Duckwater	0.0005	0.0033	0.0014	0.0028	
Ely	0.0004	0.0016	0.0009	0.0009	
Garden Valley	0.0007	0.0019	0.0013	0.0011	
Goldfield	0.0004	0.0025	0.0012	0.0013	
Henderson	0.0004	0.0029	0.0015	0.0011	
Indian Springs	0.0006	0.0027	0.0012	0.0010	
Las Vegas	0.0018	0.0046	0.0027	0.0025	
Mesquite	0.0006	0.0034	0.0017	0.0012	
Milford	0.0003	0.0033	0.0015	0.0012	
Nyala	0.0006	0.0022	0.0011	0.0009	
Overton	0.0007	0.0026	0.0016	0.0015	
Pahrump	0.0004	0.0021	0.0013	0.0014	

Table 1. Gross Alpha Analytical Results for the First Quarter of Calendar Year 2007	
(Average analytical error, +/- 0.0007)	

Pioche	0.0004	0.0033	0.0014	0.0010
Rachel	0.0007	0.0014	0.0010	0.0011
Sarcobatus	0.0006	0.0053	0.0021	0.0017
Shoshone	0.0005	0.0024	0.0012	0.0013
St. George	0.0005	0.0023	0.0016	0.0012
Stone Cabin	0.0003	0.0014	0.0010	0.0010
Tonopah	0.0005	0.0022	0.0012	0.0011
Twin Springs	0.0004	0.0020	0.0012	0.0010

Station	Minimum	Maximum	Average	2006 Average
	(pCi/m <sup>3</sup> )	$(pCi/m^3)$	(pCi/m <sup>3</sup> )	(pCi/m <sup>3</sup> )
Alamo	0.011	0.027	0.021	0.022
Amargosa	0.011	0.030	0.020	0.022
Beatty	0.010	0.035	0.023	0.022
Boulder City	0.011	0.042	0.026	0.025
Caliente	0.011	0.038	0.024	0.024
Cedar City	0.014	0.031	0.020	0.020
Delta	0.012	0.069	0.030	0.022
Duckwater	0.012	0.029	0.019	0.028
Ely	0.007	0.025	0.017	0.020
Garden Valley	0.014	0.029	0.021	0.022
Goldfield	0.009	0.030	0.020	0.020
Henderson	0.011	0.035	0.023	0.022
Indian Springs	0.013	0.035	0.021	0.020
Las Vegas	0.014	0.045	0.025	0.025
Mesquite	0.015	0.047	0.025	0.024
Milford	0.014	0.056	0.030	0.024
Nyala	0.011	0.035	0.019	0.020
Overton	0.013	0.035	0.025	0.024
Pahrump	0.014	0.032	0.023	0.022

Table 2. Gross Beta Analytical Results for the First Quarter of Calendar Year 2007.(Average analytical error, +/- 0.003)

Pioche	0.010	0.029	0.019	0.020
Rachel	0.011	0.032	0.021	0.022
Sarcobatus	0.013	0.039	0.023	0.023
Shoshone	0.013	0.042	0.025	0.024
St. George	0.016	0.042	0.027	0.024
Stone Cabin	0.013	0.025	0.018	0.020
Tonopah	0.012	0.028	0.019	0.020
Twin Springs	0.012	0.039	0.022	0.021

Station	Cs-137 (pCi/sample)	Cs-137 (MDC)	Be-7 (pCi/m <sup>3</sup> )	Pb-210 (pCi/m <sup>3</sup> )
Alamo	0.0	8.2	0.024	0.025
Amargosa	0.9	9.0	0.109	0.010
Beatty	0.0	16.0	0.094	N.D.
Boulder City	0.2	8.1	0.034	0.021
Caliente	0.4	9.1	0.033	0.018
Cedar City	-1.1	11.0	0.037	0.018
Delta	0.9	10.0	0.030	N.D.
Duckwater	-0.1	8.6	0.032	N.D.
Ely	0.2	11.0	0.033	N.D.
Garden Valley	0.4	10.0	0.042	0.018
Goldfield	-1.1	11.0	0.111	0.018
Henderson	0.0	8.3	0.038	N.D.
Indian Springs	0.0	13.0	0.118	N.D
Las Vegas	0.0	12.0	0.035	0.016
Mesquite	3.7	11.0	0.038	0.016
Milford	1.6	8.7	0.037	0.027
Nyala	-0.1	11.0	0.032	0.011
Overton	0.0	14.0	0.030	N.D.
Pahrump	0.1	10.0	0.071	0.020

Table 3. Gamma Spectroscopy Results for the First Quarter of Calendar Year 2007.

Pioche	0.8	7.6	0.041	N.D.
Rachel	1.9	9.2	0.090	0.019
Sarcobatus	-1.0	8.5	0.087	0.016
Shoshone	2.1	6.0	0.059	N.D.
St. George	-1.5	11.0	0.038	0.016
Stone Cabin	-1.0	11.0	0.042	0.013
Tonopah	-0.3	8.8	0.099	N.D.
Twin Springs	1.1	8.6	0.026	N.D.

MDC Be-7 =  $0.022 \text{ pCi/m}^3$  Pb-210 =  $0.006 \text{ pCi/m}^3$  N.D. = not detected

Station	First Quarter Exposure (mR)	Est. Annual Exposure (mR/yr)	2006 TLD Exposure (mR/yr)	2006 PIC Exposure (mR/yr)
Alamo	28	112	121	117
Amargosa	25	109	109	111
Beatty	37	149	142	137
Boulder City	25	94	108	132
Caliente	stolen			
Cedar City	25	100	96	99
Delta	26	104	101	94
Duckwater	30	120	N.A.	126
Ely	25	110	115	98
Garden Valley	37	163	152	162
Goldfield	30	127	125	133
Henderson	30	113	145	132
Indian Springs	25	109	98	102
Las Vegas	24	90	103	93
Medlins Ranch	32	138	142	146
Mesquite	26	98	107	103
Milford	36	145	144	183
Nyala	27	116	112	117
Overton	23	87	95	89
Pahrump	19	83	78	72

Table 4.TLD Analytical Results for the First Quarter of Calendar Year 2007

Pioche	28	112	113	130
Rachel	32	128	134	137
Sarcobatus	39	157	154	154
Shoshone	29	126	113	132
St. George	23	92	89	82
Stone Cabin	32	138	140	151
Tonopah	32	128	132	140
Twin Springs	39	170	153	171